

MULTIPLE PHASE CROSS-LINKED COMPOSITIONS AND USES THEREOF

CROSS-REFERENCE TO RELATED APPLICATION

Priority under 35 U.S.C. § 119(e) is claimed to Provisional Application Serial No. 60/212,511, filed June 19, 2000, incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

The present invention relates to materials, methods for their preparation, and compositions including pharmaceutical compositions that comprise a cross-linked matrix comprising a polymer and multiple phases. Such multiple-phase matrices or compositions exhibit new and useful physical properties including stability of oil-water emulsions, and controlled release kinetic profiles of active agents contained therein, making them suitable for controlled release formulations of various agents such as therapeutic agents for uses including the prophylaxis or treatment of conditions and diseases.

BACKGROUND OF THE INVENTION

Therapeutic agents with short half lives, such as most proteins, must be administered by injection at closely repeated intervals to maintain therapeutic benefit, since their in vivo half-lives are minutes to hours. A prominent approach for extending the half-life of a protein to a period of hours or days is to covalently append to it one or more chains of poly(ethylene glycol) (PEG). Appended PEG chains may provide the favorable pharmacologic properties of protection of the

1 underlying protein from immune surveillance and proteolytic enzymes, in addition to the lower
2 rate of clearance from the bloodstream (Davis, S., Abuchowski, A., Park, Y. K. and Davis, F. F.
3 (1981) Clin. Exp. Immunol. 46, 649-652.). However, the successful use of this "pegylation"
4 technology is highly and unpredictably dependent on both the particular protein and the
5 conjugation chemistry, and is effective for a few days at most. It is also not directly suited to all
6 short-lived therapeutic agents.

7
8 Another approach to extending the in vivo lifetime of a therapeutic agent is to administer that
agent encapsulated in a sustained release depot. Protein encapsulation processes that require the
use of organic solvents or heating potentially physically modify, i.e. denature, a protein drug. A
process for preparing protein microparticles by heating in the presence of polymers is described
by Woiszwilllo et al. (U. S. Patent 5,849,884). A process in which the protein drug is contacted
with an organic solvent is described by Zale et al. (U.S. Patent 5,716,644).

ENCLOSURE
16 Encapsulation processes that require chemical bond formation among the encapsulation reagents
might have reactions that unintentionally chemically modify the protein. Thus, this latter method
17 is less favored, since for the example of proteins, which are typically composed of amino acids
18 having a variety of side chain functional groups, chemical modification may impair the
19 pharmacological activity. The same impairment may be imparted to other therapeutic agents.

20
21 It is toward the development of new controlled release delivery systems for small-molecule
22 drugs, proteins and other therapeutic agents, particularly for those with short in-vivo lifetimes,
23 that the present application is directed. Furthermore, the new and useful properties of such

1 controlled release delivery materials have found uses beyond the pharmaceutical uses, in the
2 handling, storage, and delivery of industrial agents.

3
4 The citation of any reference herein should not be construed as an admission that such reference
5 is available as "Prior Art" to the instant application.

6
7 SUMMARY OF THE INVENTION

8 In its broadest aspect, the present invention is directed to compositions and methods for
9 preparing such compositions, the compositions being cross-linked polymer matrices comprising
10 a homogeneous plurality of phases, one of which is an aqueous phase. The methods for
11 preparing such compositions comprise preparing a homogeneous mixture of at least two phases,
12 one of which is an aqueous phase, and at least one polymer capable of being cross-linked present
13 in at least one of the phases, and forming cross-links between the polymer molecules. The phase
14 other than the aqueous phase may be one or more oil (lipid) phases or one or more solid phases,
15 or multiple different combinations of the phases, such as two solid phases each comprising a
16 different agent in a single aqueous phase or in an emulsion of the aqueous and oil (lipid) phases,
17 or an emulsion of two different oil (lipid) phases in a single aqueous phase. Preferably, the
18 polymer is in the aqueous phase.

19
20 In a preferred embodiment, at least one active agent is present in at least one of the phases, such
21 that the at least one agent is physically entrapped within the composition. One or more
22 excipients may be included in the composition to aid in the formation, stability and/or release
23 characteristics of the composition, such as a surfactant to aid in the formation of an emulsion, a

1 polymeric counterion to aid in the insolubilization of a polymeric active agent within the
2 composition, or a proteinase inhibitor to maintain the stability of a proteinaceous active agent
3 within the matrix.
4

5 As noted above, an agent may be physically entrapped within one or more phases in the matrix of
6 the invention. Such physical entrapment generally relates to and refers to the cross-linking of the
7 polymer which non-covalently entraps the components of the composition, including any
8 suspended (solid phase) material, the emulsion, or any additional phases present. The active
9 agent need not necessarily be present in the same phase that the polymer is present, which as
10 noted above is preferably the aqueous phase. A preferred embodiment comprises an aqueous and
11 an oil (lipid) phase, with the polymer in the aqueous phase and the therapeutic agent entrapped
12 within the polymer within the oil (lipid) phase or the aqueous phase.
13

14 In one embodiment, the at least one active agent is a therapeutic agent. The therapeutic agent
15 may be a small organic molecule, nucleic acid, peptide, polypeptide, protein, carbohydrate,
16 vaccine, adjuvant, lipid, or it may be a virus or cell, although it is not limited to any particular
17 compound, biomolecule or entity. Such compositions have desirable controlled release
18 properties such that an entrapped therapeutic agent or agents is released from the matrix under
19 zero order, pseudo zero order or first order kinetics. The release characteristics are adjustable by
20 selection of the appropriate phases, polymer(s), cross-linking agent(s), and excipients, among
21 other factors.
22

23 The compositions of the invention may be prepared from a mixture of at least two phases, one of

1 which is an aqueous phase and at least one of which comprises at least one therapeutic agent, and
2 a polymer capable of being cross-linked, and forming cross-links between the polymer molecules
3 to form a cross-linked matrix entrapping the at least one therapeutic agent. The cross-linking can
4 be performed before, during, or after the matrix is administered to the animal. For example, the
5 cross-linking reaction can be initiated in vitro, and the mixture, while undergoing cross-linking,
6 may be injected into a bodily compartment of an animal, wherein the injected bolus continues to
7 cross-link and harden in situ. In another embodiment, a cross-linked matrix after formation can
8 be implanted or inserted into the location of the body from which delivery of the agent is desired.
9 The compositions may also be introduced at either end of the gastrointestinal tract for
10 transmucosal absorption.

11
12
13
14
15
16 The additional one or more phases other than the aqueous phase may be an oil (lipid) phase, or a
17 solid phase. The oil (lipid) phase is preferably a compound or mixture thereof which is a liquid
18 at the temperature at which the compositions of the invention are used, for example, for sustained
19 release in the body or in an industrial setting. Non-limiting examples of suitable oil or lipid
20 phase components include fatty acid esters, such as lower alcohol esters of myristic acid, high
21 molecular weight fatty acids, and oils such as food oils, by way of illustration. The solid phase
22 may be a compound or agent which is insoluble in the aqueous phase. It may also be a
23 preformulated solid component, such as a microsphere or microfiber; in the case of microspheres,
another phase, such as an aqueous or lipid phase, may be present within the solid microsphere.

24
25
26 The invention is also directed to a method for the controlled release of at least one therapeutic
agent by administering to a site in the body a composition of the invention as described above.

1 The controlled release of the at least one therapeutic agent from the pharmaceutical composition
2 of this aspect of the invention may occur as a consequence of diffusion from the at least one
3 phase of the matrix wherein the active agent resides, or biodegradation of the matrix by an
4 *in-vivo* degradation pathway such as via reducing agents, reductases, S-transferases, peptidases,
5 proteases, non-enzymatic hydrolysis, esterases or thioesterases. As will be seen below, a
6 remarkable and surprising finding herein is that the presence of multiple phases beneficially
7 influences the controlled release characteristics of an active agent in the composition, whether or
8 not the active agent is contained within any particular additional phase. The release may be zero
9 order, pseudo-zero order or first order. Moreover, the ratio among the aforementioned types of
10 stable and labile cross-linking bonds, among other factors, may be used to regulate the
11 persistence of the composition within the body and the release kinetics of the entrapped
12 therapeutic agents. For example, a ratio of thioether, thioester and disulfide bonds may provide
13 the proper release pharmacokinetics for a composition of the invention placed in a particular
14 bodily site that is exposed to esterases as well as reducing activity.

15 The mixture may comprise one or more excipients that modulate one or more properties of the
16 cross-linked matrix, such as swelling of the polymer, diffusion or partitioning of the therapeutic
17 agent, or formation or maintenance of an emulsion. Such excipients include, by way of non-
18 limiting example, mono- or divalent metal ions, anions or ionic polymers, proteins such as serum
19 albumin, surfactants and polymers such as dextran. Moreover, components may be added to the
20 composition to provide enhanced stability of any therapeutic agents contained therein, for
21 example, proteinase inhibitors to maintain the stability of entrapped proteinaceous therapeutic
22 agents. Such inhibitors may be present in the aqueous, lipid or solid phase, for example, in the
23

1 form of microspheres. Slow release of the proteinase inhibitor from the microsphere protects the
2 entrapped protein from attack by proteinases from the environment of the composition (such as
3 one implanted in the body) from attaching the therapeutic agent.

4
5 The polymer of the materials and compositions of the invention comprises at least two functional
6 or reactive groups which may particulate in cross-linking to form the matrix entrapping the
7 agent, and may be a homopolymer or a copolymer. Any of a large number of such polymers or
8 combinations may be used. The polymer may have a backbone such as but not limited to a
9 polyalkylene oxide such as poly(ethylene glycol) (PEG or poly[ethylene oxide]),
10 carboxymethylcellulose, dextran, modified dextran, polyvinyl alcohol,
11 N-(2-hydroxypropyl)methacrylamide, polyvinyl pyrrolidone, poly-1,3-dioxolane,
12 poly-1,3,6-trioxane, polypropylene oxide, a copolymer of ethylene/maleic anhydride, a
13 polylactide/polyglycolide copolymer, a polyaminoacid, a copolymer of poly(ethylene glycol) and
14 an amino acid, and a polypropylene oxide/ethylene oxide copolymer. Poly(ethylene glycol) is
15 preferred. The foregoing polymer or polymers used to form the cross-linked matrix may
16 independently have one or more types of functional groups which serve as sites for cross-linking.
17 Such functional groups may be amino groups, carboxyl groups, thiol groups, and hydroxyl
18 groups, by way of non-limiting examples. By way of example, the polymer may be derived from
19 a poly(ethylene glycol) (PEG) derivative such as but not limited to α,ω -dihydroxy-PEG and
20 α,ω -diamino-PEG, which may be cross-linked via hydroxy or amino groups. Another polymer
21 with thiol functional groups may be prepared from, for example, α,ω -diamino-poly(ethylene
22 glycol) and thiomalic acid; α,ω -dihydroxy-poly(ethylene glycol) and thiomalic acid; or
23 α,ω -dicarboxy-PEG subunits and lysine, wherein free carboxy groups on the lysine residue are

1 derivatized to provide thiol groups. In a particular embodiment, the poly(ethylene glycol)
2 subunit size is from about 200 to about 20,000 Da. In a more preferred embodiment, the
3 poly(ethylene glycol) subunit size is from about 600 to about 5,000 Da.

4
5 Preferably, the polymer comprises at least two thiol groups, and may be a homopolymer or a
6 copolymer.

7
8 Such moieties may be cross-linked by reagents capable of forming covalent bonds between the
9 functional groups, such as but not limited to homobifunctional and heterobifunctional cross-
10 linking agents. A preferred moiety is a thiol group, and a preferred cross-linking agent is one
11 that forms thioether bonds, such as a vinylsulfone or maleimide, but the invention is not so
12 limiting. Other cross-linking reagents, such as a pyridyldithio-containing reagent, or oxidation,
13 may be used to generate reducible cross-links. Combinations of cross-linking reagents may be
14 used, as mentioned above, to provide a ratio of cross-link types which generate the desired
15 release characteristics of the composition. The preferred thiol-containing polymer may have
16 from 2 to about 20 thiol groups. Preferably, the polymer has from about 3 to about 20 thiol
17 groups, and most preferably, the polymer has from about 3 to about 8 thiol groups. In one
18 embodiment, the thiol groups on the polymer are sterically hindered.

19
20 As noted above, the release rate of the therapeutic or other agent in the composition of the
21 invention may be regulated by the biodegradability of the cross-linked polymer matrix. As
22 multiply types of polymers and/or multiple types of cross-links may be formed, the degradation
23 rate may be adjusted by varying the ratio or types of cross-links, and the stability or lability

1 thereof, in the composition. For example, the ratio of reducing agent-sensitive disulfide bonds,
2 esterase-sensitive ester bonds, and stable thioether bonds may be selected to provide the desired
3 release kinetics of one or more entrapped agents.

4
5 As mentioned above, any of various conditions and/or reagents may be used to effect the cross-
6 linking of the polymer, depending on the particular functional groups on the polymer. By way of
7 non-limiting example, the conditions that cause cross-linking of the thiol groups on a thiol-
8 containing polymer may be reaction in the presence of an oxidizing agent or reaction with a
cross-linking agent. In the aspect of oxidation, the oxidizing agent may be by way of non-
limiting example, molecular oxygen, hydrogen peroxide, dimethylsulfoxide, and molecular
iodine. In the aspect where a cross-linking agent is used, the cross-linking agent may be a
bifunctional disulfide-forming cross-linking agent or a bifunctional thioether-forming
cross-linking agent. In a preferred embodiment, the cross-linking agent is a long-chain
cross-linking agent, with a molecular weight of about 300 to about 5,000 Da. Non-limiting
examples of suitable cross-linking agent include 1,4-di-[3',2'-pyridyldithio(propion-
16 amido)butane]; α,ω -di-O-pyridyldisulfidyl-poly(ethylene glycol); a vinyl sulfone such as
17 α,ω -divinylsulfone-poly(ethylene glycol); 1,11-bis-maleimidotetraethylene glycol; and
18 α,ω -diiodoacetamide-poly(ethylene glycol).

19
20 For other functional groups or a combination of a thiol group and another group, any appropriate
21 bifunctional cross-linking agent may be selected which will achieve the desired cross-linking of
22 the functional groups and formation of the cross-linked polymer.

1 In another aspect, the polymer additionally comprises a functional group, which may derivatized
2 for example with a label, such as a contrast/imaging agent, radionuclide, chromophore,
3 fluorophore, red or near-infrared fluorophore, or nonradioactive isotope. In another embodiment,
4 the label is a metabolically stable polymer component that after degradation of the polymer is
5 detectable in the urine. In another embodiment, the cross-linking agent used to cross-link the
6 polymer additionally comprises a functional group, such as a label.

7
8 In another related aspect, the delivering of at least one therapeutic agent to a bodily compartment
9 under controlled release conditions is provided by situating in the bodily compartment a
10 pharmaceutical composition comprising a matrix as described hereinabove. The bodily
11 compartment may be subcutaneous, oral, intravenous, intraperitoneal, intradermal, subdermal,
12 intratumor, intraocular, intravisceral, intraglandular, intravaginal, intrasinus, intraventricular,
13 intrathecal, intramuscular, or intrarectal, by way of non-limiting examples. The composition of
14 the invention may be provided to the bodily compartment by a route such as but not limited to
15 subcutaneous, oral, intravenous, intraperitoneal, intradermal, subdermal, intratumor, intraocular,
16 intravisceral, intraglandular, intravaginal, intrasinus, intraventricular, intrathecal, intramuscular,
17 or intrarectal.

18
19 In yet a further aspect, the invention is directed to a method of preparing a cross-linked hydrogel
20 drug depot, the method comprising: preparing a mixture comprising at least one therapeutic
21 agent in a plurality of phases and a polymer system capable of forming a cross-linked hydrogel
22 matrix, the polymer system comprising a first polymer having a plurality of functional groups,
23 and a second polymer or long-chain compound having two or more functional or reactive groups;

1 and forming linkages between the functional groups of the first polymer and the functional or
2 reactive groups of the second polymer so as to form a cross-linked hydrogel matrix having a
3 plurality of phases and the therapeutic agent physically entrapped therein. The plurality of
4 phases are as described hereinabove. The first and second polymer may be the same or different.
5 The first or second polymer may be a polyalkylene oxide, and either or both may be a
6 homopolymer, a copolymer or a combination thereof. They may have one or more
7 biodegradable linkages. In a preferred embodiment, one polymer comprises thiol groups and the
8 other comprises vinylsulfone or maleimide groups. Reaction of the vinylsulfone or maleimide
9 groups with the thiol groups forms cross-links. In another embodiment, the first and second
10 polymers comprise thiol groups, and a homobifunctional thiol-reactive cross-linking agent is
11 used to form cross-links. In these examples, the plurality of thiol groups may be between 2 and
12 20. The second polymer may be a long-chain cross-linking agent.

13
14 The releasing of a therapeutically effective amount of the therapeutic agent from the
15 cross-linked hydrogel matrix may occur over a time course of three or more, five or more, ten or
16 more, fifteen or more, or twenty or more days. Release of weeks to months by the compositions
17 of the invention is also embraced herein. The controlled release profile may comprise a desired
18 initial bolus release profile followed by a release profile such as but not limited to zero order,
19 pseudo zero order, and first order.

20
21 The invention is further directed to a method of administering a therapeutic agent to a mammal,
22 the method comprising: preparing a mixture comprising a hydrogel-forming polymer having two
23 or more thiol groups, a cross-linker comprising two or more vinylsulfone or maleimide groups,

1 and a therapeutic amount of drug and a plurality of phases; and injecting into a particular bodily
2 compartment of the mammal with the mixture whereby a hydrogel drug depot is formed at the
3 site of injection having said drug temporarily entrapped therein. Furthermore, the invention is
4 also directed to a method of administering a therapeutic agent to a mammal by preparing a
5 mixture comprising a hydrogel-forming polymer having two or more vinylsulfone groups, a
6 cross-linker comprising two or more thiol groups, a therapeutic amount of drug, and a plurality
7 of phases; and injecting said mammal with said mixture whereby a hydrogel drug depot is
8 formed at the site of injection having the drug temporarily entrapped therein. In either of the
9 foregoing methods, the cross-linker may comprise a hydrogel forming polymer, and may further
10 comprise releasing a therapeutically effective amount of the therapeutic agent from said hydrogel
11 drug depot over a time course of three or more days. The injecting may be subcutaneous.

12
13 In a further embodiment, the present invention is directed to a hydrogel drug depot comprising a
14 therapeutic agent physically entrapped within a polymer matrix comprising a thioether
15 cross-linked hydrogel matrix and a plurality of phases. The hydrogel matrix may comprise a
16 polyalkylene oxide, which may be a homopolymer, copolymer or combination thereof of
17 poly(ethylene glycol) or derivative thereof. The polymer matrix may comprise a controlled
18 release kinetic profile characterized by release of a therapeutically effective amount of the
19 therapeutic agent from the thioether cross-linked hydrogel matrix over a time course of three or
20 more, five or more, ten or more, fifteen or more, or twenty or more days. The controlled release
21 kinetic profile may comprise an initial bolus release profile followed by a release profile such as
22 zero order, pseudo zero order, or first order. The hydrogel depot may comprise one or more
23 excipients that modulate one or more properties of the thioether cross-linked hydrogel matrix,

1 such as diffusion, swelling, partitioning of the therapeutic agent, or formation or maintenance of
2 an emulsion. Such excipients include, by way of non-limiting example, mono- or divalent metal
3 ions, anions or ionic polymers, proteins such as serum albumin, surfactants, and polymers such
4 as dextran. A proteinase inhibitor may be used. One or more may be present in the composition.

5
6 The therapeutic agent of the hydrogel drug depot may be, by way of non-limiting example, a
7 small organic molecule, nucleic acid, peptide, polypeptide, protein, carbohydrate, vaccine,
8 adjuvant, or lipid.

9
10 The cross-linked hydrogel matrix of the hydrogel drug depot may be formed by cross-linking a
11 first polymer containing two or more thiol groups with a second polymer or long-chain
12 compound containing two or more vinyl sulfone groups. The first polymer may comprise a
13 molecular weight of 200 to 20,000 Daltons; the second polymer or long-chain compound may
14 comprise a molecular weight of 100 to 5,000 Daltons. The first polymer may comprise between
15 2 and 20 thiol groups. The first and second polymers may be in a defined molar ratio for
16 controlling the controlled release kinetic profile of the hydrogel drug depot. The thioether
17 cross-linked hydrogel matrix may comprise one or more biodegradable linkages.

18
19 In yet a further aspect, the invention is directed to a hydrogel drug depot system comprising a
20 compound of interest, a plurality of phases, a first polyalkylene oxide polymer containing two or
21 more thiol groups, a second polyalkylene oxide polymer containing two or more vinyl sulfone
22 groups that are capable of covalently bonding to one another to form a thioether cross-linked
23 hydrogel matrix, the hydrogel drug depot system having a controlled release kinetic profile

1 characterized by sustained release of the compound of interest from the thioether cross-linked
2 hydrogel matrix over a time course of three or more days and in some embodiments extending up
3 to several months. The polyalkylene oxide may be poly(ethylene glycol) or a derivative thereof;
4 the hydrogel matrix may comprise one or more biodegradable linkages, such as but not limited to
5 an ester linkage. The hydrogel drug depot may have a controlled release kinetic profile
6 comprising an initial bolus release profile followed by a release profile such as zero order,
7 pseudo zero order, or first order.

8
9 The invention is also directed to a kit for forming a hydrogel drug depot comprising an agent of
10 interest such as a therapeutic agent or a diagnostic agent, the kit including an aqueous phase, an
11 oil or lipid phase, a surfactant, a polymer with two or more functional groups, and a cross-linking
12 agent capable of forming a cross-links among the functional groups. In the use of the kit, a
13 therapeutic agent is added to the components and cross-linking induced, in accordance with one
14 of the aforementioned processes of forming the matrix in vitro, or forming it in situ by injecting
15 the components soon after mixing, such that the matrix is not yet polymerized and can pass
16 through a needle or cannula, and full cross-linking occurs in situ.

17
18 In a particular embodiment, the invention is directed to a kit for forming a hydrogel drug depot
19 comprising an agent of interest such as a therapeutic agent or a diagnostic agent, a polymer
20 having two or more thiol groups, and a low molecular weight, polymer or long-chain cross-
21 linking compound having two or more vinylsulfone groups, and a plurality of phases, wherein
22 said polymer and said cross-linker are capable of covalently bonding to one another under
23 physiological conditions to form a thioether cross-linked hydrogel matrix so as to entrap the

agent of interest therein. The hydrogel matrix may comprise polyalkylene oxide. In the foregoing kit, the polyalkylene oxide may be a homopolymer, copolymer or combination thereof of poly(ethylene glycol) or derivative thereof. The polymer matrix comprises a controlled release kinetic profile characterized by release of a therapeutically effective amount of the therapeutic agent from the thioether cross-linked hydrogel matrix over a time course of three or more, five or more, ten or more, fifteen or more, or twenty or more days. Release over weeks to months is also embodied herein. The controlled release kinetic profile may comprise an initial bolus release profile followed by a release profile such as zero order, pseudo zero order, or first order. The kit may include one or more excipients that modulate one or more properties of the thioether cross-linked hydrogel matrix, such as, but not limited to diffusion and swelling. The therapeutic agent is selected from the group consisting of small organic molecule, nucleic acid, peptide, polypeptide, protein, carbohydrate, vaccine, adjuvant, and lipid. The diagnostic agent may be a contrast/imaging agent, radionuclide, chromophore, fluorophore, red or near-infrared fluorophore, or a non-radioactive isotope. The kit may also have other types of polymers and cross-linkers.

In the aforementioned kits, the polymer may have a molecular weight of 200 to 20,000 Daltons. The cross-linking agent, whether a small molecule, polymer or long-chain compound, may have a molecular weight of 100 to 5,000 Daltons. The polymer may have between 2 and 20 thiol groups. The polymer and cross-linking agent may be provided in preformed aliquots for admixing to generate a defined molar ratio of the first and second polymers for controlling the controlled release kinetic profile of the hydrogel drug depot.

1 In another aspect of the invention, a method of producing a kit according to the above may be
2 performed by assembling in the kit an agent of interest such as a therapeutic agent or a diagnostic
3 agent, a first polymer having two or more thiol groups, and a second polymer or long-chain
4 compound having two or more vinyl sulfone groups, wherein the first polymer and the second
5 polymer or long-chain compound are capable of covalently bonding to one another under
6 physiological conditions to form a thioether cross-linked hydrogel matrix so as to entrap the
7 agent of interest therein.

8
9 These and other aspects of the present invention will be better appreciated by reference to the
10 following drawing and Detailed Description.

11 BRIEF DESCRIPTION OF THE DRAWING

12 Figure 1 shows the in vitro release of quinine sulfate monohydrate from two different
13 formulations of the thiol containing polymer hydrogel, formulation I having an aqueous phase,
14 and oil phase and a solid phase, and formulation II having an aqueous phase and a solid phase.
15

16
17 Figure 2 shows another in vitro release of quinine sulfate monohydrate from two different
18 formulations of the thiol containing polymer hydrogel.

19
20 Figure 3 shows the in-vivo release of quinine sulfate monohydrate from two different
21 formulations of the thiol containing polymer hydrogel, the Rgel formulation having a aqueous
22 phase and a solid phase, and the Egel formulation an aqueous, an oil phase and a solid phase. The
23 insert shows the initial release profiles of the two formulations.

1 Figure 4 shows the in-vitro release of salmon calcitonin from two different formulations of the
2 thiol containing polymer hydrogel of the invention.

3
4 Figure 5 shows the in-vivo release of salmon calcitonin from two different formulations of the
5 thiol containing polymer hydrogel, Rgel formulation having a single aqueous phase and Egel
6 formulation an aqueous and an oil phase.

7 8 DETAILED DESCRIPTION OF THE INVENTION

9 The present invention provides new and useful materials including compositions and
10 pharmaceutical compositions, and methods for their preparation and administration, based upon
11 the surprising and remarkable discovery by the inventors herein that cross-linking of a polymer
12 and formation of a cross-linked polymer matrix in a multiple phase system comprising an
13 aqueous phase and at least one other phase, preferably at least an oil (lipid) phase, provides a
14 material with various heretofore unknown properties useful for a variety of industrial and
15 medical applications, among others. By physically entrapping, for example, an active agent in
16 the composition in one or more of the phases, particular storage and handling features of the
17 material may be provided, and such materials may be prepared with desirable release properties
18 for the one or more entrapped agents. As will be elaborated upon below, such multiple-phase
19 systems include an aqueous phase and a solid phase, or an aqueous phase and an oil (lipid) phase,
20 or an aqueous, oil (lipid) and solid phase. Furthermore, the aqueous and oil (lipid) phases may
21 be provided in the form of an emulsion. An emulsion is a preferred multiple phase system. The
22 terms "solution," "mixture," and "suspension" are used interchangeably to refer to the
23 compositions herein comprising a plurality of phases before the matrix is cross-linked, such as a

1 suspension of solid particles in an aqueous phase or an oil-aqueous phase emulsion. Oil and lipid
2 are interchangeably used to refer to a liquid, water-insoluble phase. Various agents, other than
3 the active agent(s), herein termed excipients, may be included in the compositions to enhance the
4 formation or stability of the emulsion, to maintain the separate phases, or to modulate the
5 partitioning of an active agent among the phases and to modulate the storage (retention) or
6 release characteristics of the composition. Moreover, multiple oil phases or solid phases may be
7 present in the aqueous phase, for example, two or more types of oils or solid particles of different
8 compositions. In a preferred embodiment, an emulsion is formed from an aqueous and an oil
9 phase, with a surfactant or detergent.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2

1 formation of the cross-linked polymer. Moreover, the invention is not so limiting as to the nature
2 of the agent physically entrapped in the compositions herein.

3
4 By way of the non-limiting example of a pharmaceutically-useful agent (therapeutic agent) in the
5 instant composition, the composition may be prepared to release the agent(s) with a controlled
6 release kinetic profile in vivo, such as zero order, pseudo zero order or first order release. A
7 preferred release is a constant rate of release over time. As will be seen below, the compositions
8 are prepared using a polymer with functional and/or reactive groups that may participate in cross-
9 linking reactions, and another compound, such as but not limited to a polymer, which reacts with
10 and cross-links the polymer with functional or reactive groups. A mixture of the multiple phases,
11 polymer, and optional active agent(s) and excipient(s) is prepared, and then cross-linking of the
12 polymer is initiated by placing the mixture under conditions which cause cross-linking, such as
13 exposure to a cross-linking agent, heating, cooling, polymerization-inducing radiation, etc.
14 Moreover, the compositions may be prepared such that the material may be readily deposited in a
15 bodily compartment without the need for surgery, by injecting through a needle or cannula the
16 composition of the invention while in liquid form, the cross-linking reaction solidifying the
17 composition in situ.

18
19 The compositions of the invention comprise a polymer matrix prepared from polymers bearing
20 moieties, such as thiol moieties, which are capable of being cross-linked by any of a number of
21 processes, such as oxidation or by use of a bifunctional cross-linking agent, to physically entrap
22 the therapeutic agent within the cross-linked polymer. The matrices are prepared by cross-
23 linking the polymer in the presence of the therapeutic agent(s), such that entrapment occurs

1 during cross-linking. The invention is directed to compositions and pharmaceutical
2 compositions prepared by these methods, methods of preparing the compositions by cross-
3 linking the polymer to entrap the agent therein, as well as to methods for administering the
4 composition to an animal, for instance, by injecting the composition of the invention into an
5 animal during the process of cross-linking such that the mixture is liquid or semi-solid during
6 injection, but soon after injection completes the cross-linking process and forms the matrix
7 (depot) with the aforementioned release characteristics. Thus, the cross-linking of the polymer
8 may be performed during the manufacture of the composition, which is subsequently
9 administered to or implanted at the desired site; in another embodiment, a mixture of the
10 therapeutic agent(s), the polymer and the cross-linking agent is administered to the desired site at
11 the time of or just after initiation of the cross-linking reaction, such that the mixture can be
12 readily deposited at the desired site, and the cross-linking subsequently occurs or is completed in
13 the bodily compartment to form the matrix. In all of the foregoing examples, the agent or agents
14 and multiple phases are physically entrapped within the cross-linked polymer.

15 The term agent or active agent refers to the substance for which the matrix of the invention may
16 be used to hold, deliver, stabilize, release, carry, transport, store, or otherwise handle for any
17 purpose for which the agent may be used. As noted herein, a preferred agent is a therapeutic
18 agent, but the agent may be any agent. The term therapeutic agent should not be considered
19 limiting to medically useful agents. The composition of the invention may not comprise any
20 active agent, the cross-linked multiphase composition having useful properties itself.

21
22
23 As used herein, a phase refers to a distinct liquid or solid phase, such as an aqueous, solid, or oil

phase, and as will be seen below, a composition or matrix of the invention comprises two or more phases. One of the phase is an aqueous phase. The agent may be present in one or more phases. For example, a matrix comprising an poorly water-soluble agent may comprise a solid phase (the agent), and an aqueous phase. A matrix may comprise an emulsion of an aqueous phase and an oil phase, the agent present in the aqueous phase, the oil phase, or both phases. A matrix comprising three phases may comprise an emulsion with a solid phase, the solid phase present in the aqueous phase, the oil phase, or in both. Moreover, multiple agents may be present in the compositions of the invention; for example, multiple agents suspended in an aqueous phase; multiple oil-soluble agents present in single or multiple oil phases, such as by the mixture of two emulsions, each prepared from a different oil-soluble agent, before cross-linking of the polymer. As noted above, an excipient may be used to enhance or assist in the formation of the multiple phase system, for example, by use of a surfactant such as a detergent to form the emulsion; the use of a monovalent or polyvalent metal ion or polymer to aid in the insolubilization of the active agent, or an excipient to alter the pH or other properties to partition the active agent in one or more phases and regulate or modulate its release from the one or more phases to the desired outside environment in which the composition of the invention resides. It may also include a proteinase inhibitor which prevents degradation of a proteinaceous agent within the matrix. The term excipient is used herein to refer to any compound, substance, agent, material, etc., which is not the active agent whose release is provided by the instant compositions, but agents which regulate or modulate the release, including formation of the emulsion or the matrix in general. The desired release may be no release. The excipient may be retained in the composition during the erosion or diffusion of the active agent from the composition or it may be co-released along with the active agent including in a complex with the

1 agent, but permits the active agent to have its desired activity or function after release from the
2 instant composition.

3
4 The phases present in the cross-linked matrix include an aqueous phase and at least one
5 additional phase. In a preferred embodiment, the additional phase is an oil phase, such as ethyl
6 myristate as described in the examples. Other choices of oils for the oil phase are one or more
7 compounds which are liquid at the temperature at which the compositions of the invention are
8 used, for example, for sustained release in the body or in an industrial setting. Non-limiting
9 examples of suitable oil or lipid phase components include fatty acid esters, such as lower
10 alcohol esters of caproic acid (C6), caprylic acid (C8), capric acid (C10), undecanoic acid (C11),
11 lauric acid (C12), tridecanoic acid (C13), myristic acid (C14), and palmitic acid (C16). Non-
12 limiting examples of such esters include but is not limited to caproic acid ethyl ester, caprylic
13 acid ethyl ester, capric acid ethyl ester, undecanoic acid ethyl ester, lauric acid ethyl ester,
14 tridecanoic acid ethyl ester, myristic acid ethyl ester, and palmitic acid ethyl ester. Other choices
15 for the oil phase include triglycerides that are liquid at room temperature, such as triacetin (C2),
16 tributyrin (C4), tricaproin (C6), and tricaprylin (C8). Also, fatty alcohols which are liquid at
17 room temp may be used, such as 1-octanol (C8) and 1-decanol (C10). Other examples include
18 unsaturated fatty acids such as cis-11,14-eicosadienoic acid, and unsaturated fatty acid esters
19 such as cis-11,14-eicosadienoic acid ethyl ester. Food oils such as the vegetable oils corn oil,
20 olive oil, safflower oil, and canola oil may be used. There are merely illustrative of various
21 water-immiscible liquids that may be used as an oil phase of the compositions of the invention,
22 and that may be prepared as an emulsion in combination with an aqueous phase in one
23 embodiment of the invention.

1
2 Inclusion of a surfactant such as sodium dodecyl sulfate (SDS) in the aqueous phase provides for
3 the rapid formation of an emulsion of the aqueous and oil phases. Multiple different oil phases
4 may be present. In another embodiment, a solid phase, such as a water-insoluble therapeutic
5 agent, may be present in the aqueous phase or in the emulsion. The oil phase may be any
6 water-immiscible liquid. Low molecular weight alcohol esters of fatty acids are preferred oil
7 components, but the invention is not so limited.

8
9 In another embodiment, compositions are described which comprise a cross-linked polymer
10 matrix entrapping at least one therapeutic agent, the matrix comprising a plurality of phases, at
11 least one being an aqueous phase. Methods for preparing the latter compositions are also
12 described. Thus, in this aspect, solid particles of a poorly water soluble therapeutic agent may
13 be suspended in an aqueous phase, the foregoing entrapped within the matrix. Continuous
14 release from the matrix of the molecules of the therapeutic agent that have become dissolved in
15 the aqueous phase will result in continuous solubilization of the suspended therapeutic agent into
16 the aqueous phase, thus replenishing the therapeutic agent in the aqueous phase. In a similar
17 manner, a bi-phasic system comprising an oil-water emulsion wherein the therapeutic agent is
18 present in the oil phase or aqueous phase, either dissolved or suspended (based upon its
19 solubility), also represents a controlled release system in which, for example, an oil-soluble
20 therapeutic agent with limited water solubility is entrapped in the matrix; release of the agent
21 from the aqueous phase permits redistribution of the agent from the oil phase into the aqueous
22 phase. As will be noted in more detail below, various excipients may be included in the
23 compositions herein to aid in the formation and/or stability of the composition with multiple

1 phases, particularly emulsions, as well as regulate the partitioning of the agent among the phases,
2 which further modulate the release characteristics and kinetics of the compositions.

3
4 The polymer which is cross-linked to entrap the therapeutic agents may be any cross-linkable
5 polymer, which bears two or more functional or reactive groups capable of participating in a
6 cross-linking reaction to form a matrix of the invention. Such functional groups include but are
7 not limited to amino, carboxyl, thiol and hydroxyl groups, or combinations thereof; reactive
8 groups include vinylsulfone, maleimide, pyridyldithio, and other moieties capable of reacting
9 with the aforementioned functional groups, among others. A preferred polymer is one on which
10 at least two thiol groups are present and is cross-linked with a thiol-reactive bifunctional cross-
11 linking reagent in the presence of the therapeutic agent, thus forming a cross-linked polymer with
12 the therapeutic agent physically entrapped therein. Selection of the appropriate polymer, the
13 concentration in the matrix, the extent of functional groups capable of participating in cross-
14 linking, the type of cross-linking agent, and the extent of cross-linking, and other factors will be
15 governed by such factors as the amount of therapeutic agent present in the composition, the
16 number of phases and the relative amounts of the phases including the phase in which the
17 polymer is present, in order to achieve the desired controlled release properties of the
18 composition, or retention of the active agent within the composition. In a preferred embodiment
19 for pharmaceutical agents, and in particular for proteins, the cross-linking is accomplished by
20 using sulfur chemistry for cross-linking the polymer, thereby avoiding reaction of virtually all
21 amino acid and carbohydrate side chains of, for example, a protein therapeutic agent undergoing
22 entrapment in the matrix. Although sulfur chemistry is the basis of the cross-linking preferably
23 used in this invention, disulfide bonds already present in a particular protein would be non-

1 reactive under the cross-linking conditions. Also, the sulfur atom in the thioether side chain of
2 methionine residues in the protein drug would be nonreactive. Delivery of small-molecule
3 drugs, peptides, proteins, polysaccharides, and polynucleotides including antisense nucleotides
4 are achievable using the methods described herein. Proteins containing free thiol groups
5 (cysteine residues that are not in disulfide linkage), might not be suitable for use in their native
6 form in this invention, and may need to be derivatized or otherwise protected during the
7 entrapment process. Similar considerations are given to other non-protein therapeutic agents
8 which are used in the present invention.

9
10 One advantage to using sulfur chemistry in general, and reducible cross-links in particular as
11 may be produced from oxidation of the thiol groups on the polymer or by use of a reducible
12 bond-forming cross-linking agent such as one containing a pyridyldithio (pyridyldisulfidyl)
13 group is that a cross-linked matrix formed in situ in a bodily compartment or other relatively
14 inaccessible area may be readily and facily removed by exposing the cross-linked composition
15 in situ to a reducing agent, whereupon the cross-links are broken and the composition can be
16 flushed or extracted from the site. This may be achieved, for example, when an implanted
17 release composition has achieved its desired goal of controlled releasing a therapeutic agent over
18 time, or for early removal of a device. Of course, since any remaining therapeutic agent
19 entrapped within an implanted device will be subject to rapid release when the cross-linked
20 polymer is rapidly depolymerized, considerations must be given to remove the device from the
21 site to avoid an unwanted bolus release.

22
23 However, the invention is not so limiting to sulfur chemistry to form the cross-linked matrix, and

1 polymers and cross-linking agents which achieve the desired properties may be achieved using
2 other functional and reactive groups, including both polymeric and non-polymeric cross-linking
3 agents.

4
5 With regard to pharmaceutically-useful active agents, for long term therapy (days, weeks or
6 months) and/or to maintain the highest possible drug concentration at a particular location in the
7 body, the present invention provides a sustained release depot formulation with the following
8 preferred but non-limiting characteristics: (1) the process used to prepare the matrix does not
9 chemically or physically damage the therapeutic agent, in particular proteins, thereby avoiding
10 protein inactivation or rendering the protein immunogenic; (2) the matrix maintains the stability
11 of a therapeutic agent against denaturation or other metabolic conversion by protection within the
12 matrix until release, which is important for very long sustained release; (3) the entrapped
13 therapeutic agent is released from the depot at a substantially uniform rate, following a kinetic
14 profile, and furthermore, a particular therapeutic agent can be prepared with two or more kinetic
15 profiles, for example, to provide a loading dose and then a sustained release dose; (4) the desired
16 release profile can be selected by varying the components and the process by which the matrix is
17 prepared; and (5) the matrix is nontoxic and degradable.

18
19 In the preferred but non-limiting embodiment, the cross-linked matrix of the present invention
20 entrapping at least one therapeutic agent is prepared by cross-linking a polymer for example on
21 which at least two thiol groups are present, by any one of various means, in the presence of the
22 therapeutic agent to be physically entrapped. Various polymers on which at least two thiol
23 groups are present are suitable for the use herein. The polymer on which at least two thiol

groups are present may be prepared, for example, by the reaction or derivatization of a particular polymer that does not contain thiol groups, with a thiol-containing compound, or a compound to which thiol moieties may be attached. A polymer may be prepared which has reactive terminal ends or functional groups on the ends of the polymer chain which may be subsequently derivatized to attach thiol groups. A copolymer may be prepared with repeating or alternately repeating thiol groups or functional groups which may be subsequently derivatized to have thiol groups. The extent of derivatization to provide thiol groups may be tailored to the requirements of the matrix to be formed. The foregoing examples of the types of suitable polymers is not intended to be limiting, but to be illustrative of the varieties of polymers and polymer derivatives that may be used in the practice of the invention.

In the case of thiol groups, to participate in cross-linking, the polymer has at least two thiol groups to participate in the formation of cross-links. For example, the polymer on which at least two thiol groups are present may have from 2 to about 20 thiol moieties. In a preferred embodiment, the polymer has from 3 to about 20 thiol moieties, and in a most preferred embodiment, the thiol containing polymer has from 3 to about 8 thiol moieties. These numbers of functional groups on the polymer are equally applicable to other selections of functional groups, such as amino, carboxyl and hydroxy groups, by way of non-limiting examples.

Examples of suitable subunit polymers for the preparation of the polymer on which at least two thiol groups are present include both homopolymers or copolymers. By way of non-limiting example, suitable polymers, which may be chemically modified to comprise thiol groups, include polyalkylene oxides such as poly(ethylene glycol) [also known as polyethylene glycol or PEG,

polyethylene oxide or PEO], carboxymethylcellulose, dextran, polyvinyl alcohol, N-(2-hydroxypropyl)methacrylamide, polyvinyl pyrrolidone, poly-1,3-dioxolane, poly-1,3,6-trioxane, polypropylene oxide, a copolymer of ethylene/maleic anhydride, a polylactide/polyglycolide copolymer, a polyaminoacid, a copolymer of poly(ethylene glycol) and an amino acid, or a polypropylene oxide/ethylene oxide copolymer. Such polymers are then derivatized or further polymerized to introduce thiol groups; chemical modification of the polymer may be necessary as a step prior to the further derivatization to incorporate thiol groups. For example, a polymer of the present invention may be derived from a poly(ethylene glycol) (PEG) derivative, for example, α,ω -dihydroxy-PEG or α,ω -diamino-PEG, but other derivatives are embraced herein. The polymer comprising thiol groups may be, for example, a polymer of α,ω -diamino-poly(ethylene glycol) and thiomalic acid; a polymer of α,ω -dihydroxy-poly(ethylene glycol) and thiomalic acid; or a polymer of α,ω -dicarboxy-PEG subunits and lysine wherein the free carboxy groups on the lysine residues are derivatized to form thiol groups. These polymers are only examples of possible choices, as the skilled artisan will be aware of numerous alternatives. As will be noted below, the selection of the polymer, or combinations thereof, will be guided by the desired properties of the final product, particularly the duration of release of the therapeutic agent and the release kinetics. As will also be noted below, a product of the invention may comprise more than one polymer component in order to provide two or more different release characteristics. Of course, more than one therapeutic agent may be included.

In one particular embodiment, a polymer of the present invention is derived from a poly(ethylene glycol) (PEG) derivative, for example, α,ω -dihydroxy-PEG or α,ω -diamino-PEG, but other

1 derivatives are embraced herein. Examples of such polymers with particular molecular weights
2 include α,ω -dihydroxy-PEG_{3,400} ; α,ω -dihydroxy-PEG_{1,000} ; α,ω -diamino-PEG_{3,400}; and
3 α,ω -diamino-PEG_{1,000}. PEG is known to be a particularly nontoxic polymer. These derivatized
4 PEG subunit polymers may be used as amino- and hydroxy-containing polymers for cross-
5 linking, or may be further derivatized, for example, to prepare the polymer on which at least two
6 thiol groups are present by derivatization with thiomalic acid. Thiomalic acid (also known as
7 mercaptosuccinic acid) may be replaced by dimercaptosuccinic acid, thereby doubling the
8 number of sites available for cross-linking. Increasing the extent of cross-linking the matrix
9 results in a gel with smaller pores.

10
11
12
13 As will be shown in an example below, to prepare the polymer on which at least two thiol groups
14 are present from these reactants, the thiol group of thiomalic acid is first protected by reaction
15 with trityl chloride, to produce trityl-thiomalic acid. Subsequently, the polymer on which at least
16 two thiol groups are present is prepared from the trityl-thiomalic acid and, for example,
17 α,ω -dihydroxy-PEG. Under suitable conditions, a carbodiimide is used to condense the
18 α,ω -dihydroxy-PEG with the protected thiomalic acid. After condensation, the trityl group is
19 removed by treatment with trifluoroacetic acid (TFA).

20
21 In another example, a polymer of α,ω -dicarboxy-PEG and lysine may be prepared, and
22 subsequently the free carboxy groups on the lysine residues are derivatized to form thiol groups.
23 These examples are provided by way of illustration only and such methods for adding thiol
groups to a polymer are known to those skilled in the art.

1 In a preferred embodiment using PEG as the subunit for preparing the polymer on which at least
2 two thiol groups are present, the poly(ethylene glycol) subunit size for the polymer may be from
3 about 200 to about 20,000 Da; preferably, the subunit size is from about 600 to about 5,000 Da.
4 As mentioned above, the polymer of the present invention has from 2 to about 20 thiol groups;
5 preferably from about 3 to about 20 thiol groups, and most preferably, from about 3 to about 8
6 thiol groups.

7
8 The thiol groups on the polymer on which at least two thiol groups are present may be sterically
9 hindered. It has been found that a polymer on which at least two thiol groups are present with
10 sterically hindered thiol groups tends to be nonreactive with disulfide bonds in the therapeutic
11 agent, particularly a protein, and thus does not interfere with the intramolecular disulfide bonds
12 in the protein. Furthermore, steric hindrance governs the rate at which reductive cleavage of the
13 polymer occurs in vivo. Thus, for the entrapment of proteins or other therapeutic agents with
14 disulfide bonds, a polymer on which at least two thiol groups are present, sterically hindered
15 thiol groups may be preferred. Such sterically hindered thiol groups are also preferred when
16 increased resistance to reductive cleavage is desired, for example in a longer controlled release
17 formulation. Based on the knowledge of the therapeutic agent and the particular controlled
18 release characteristics desired at the site of administration of the matrix, the skilled artisan will be
19 able to design a matrix with the desired characteristics. Examples of such sterically hindered
20 thiol groups include thiomalate, as used in the above example.

21
22 A matrix of the present invention may be prepared by cross-linking the polymer on which at least
23 two thiol groups are present in the presence of the therapeutic agent. The cross-linking of the

1 polymer on which at least two thiol groups are present may include disulfide bonds, thioether
2 bonds, and combinations thereof. Other means of covalent bond formation of thiol groups in the
3 thiol-containing polymer to effect cross-linking will be known to the skilled artisan and are
4 considered within the scope and spirit of this invention.

5
6 In one example, reaction of the polymer on which at least two thiol groups are present in the
7 presence of an oxidizing agent forms disulfide cross-links. This may be achieved by molecular
8 oxygen, hydrogen peroxide, dimethyl sulfoxide (DMSO), or molecular iodine. In other
9 embodiments, the cross-linking may be carried out by reaction with a bifunctional disulfide-
10 forming cross-linking agent, or reaction with a bifunctional thioether-forming cross-linking
11 agent. Such cross-linking agents may have a molecular weight of about 300 to about 5,000 Da,
12 and may be a polymeric cross-linking agent.

13
14 For example, the PEG-thiomalate polymer described above may be cross-linked with the non-
15 polymeric cross-linking agent 1,4-di-[3',2'-pyridyldithio(propionamido)-butane]. Alternatively,
16 a polymeric cross-linking agent such as α,ω -di-O-pyridyldisulfidyl-poly(ethylene glycol);
17 α,ω -divinylsulfone-poly(ethylene glycol); or α,ω -diiodoacetamide-poly(ethylene glycol) may be
18 used. Another thioether-forming thiol group crosslinker is 1,11-bis-maleimidotetraethylene
19 glycol, abbreviated BM(EG)₄ or BM[PEO]₄, available from Pierce. Examples of the cross-
20 linking reaction are provided in the examples below; the skilled artisan will be aware of
21 numerous other agents capable of forming the suitable matrix. As noted above, the selection of
22 the cross-linking agent is guided by the desired characteristics of the matrix product, i.e., the
23 controlled release kinetic profile and the duration of release. These factors, as well as the

1 potential reactivity of the cross-linking agent with reactive moieties on the therapeutic agent,
2 must be taken into consideration in selecting the appropriate polymer, and cross-linking agent in
3 the preparation of the product. And as mentioned above, the presence of reducible cross-links,
4 such as derived using a pyridyldithio cross-linker or oxidation, may be useful in whole or in part
5 for regulating the release characteristics of the composition, or for depolymerizing the
6 composition for removal after use.

7
8 The therapeutic agent physically entrapped in the matrix of the present invention is a compound
9 capable of being entrapped and then released in a controlled manner from the matrix. A wide
10 variety of both high molecular weight and low molecular weight compounds are suitable, and as
11 will be noted below, a compound not suitable because of its small size may be made suitable by
12 appropriate modification by for example, polymerization or conjugation to a polymer. The
13 therapeutic agent may be a small-molecule drug, protein, peptide, polysaccharide,
14 polynucleotide, or any other compound that may be entrapped in the matrix of the present
15 invention and subjected to controlled delivery in vivo. It is noted that a further advantage of the
16 present invention is that the matrix protects the therapeutic agent from degradation or other
17 metabolic processing. The agents may be for the prophylaxis or treatment of a condition or
18 disease, or for the purpose of providing controlled delivery of any suitable agent.

19
20 For example, when polymers of the following PEG polymers are prepared with thiomalic acid,
21 and then similarly cross-linked, certain properties of the polymer are obtained. The
22 α,ω -dihydroxy-PEG_{3,400} polymer subunit is conjugated via an ester bond to the thiomalic acid,
23 and the resulting product is loosely cross-linked. Likewise, a loosely cross-linked product is

1 formed from thiomalic acid and α,ω -diamino-PEG_{3,400}, the thiomalic acid linked through an
2 amide bond to the PEG subunit. In contrast, α,ω -dihydroxy-PEG_{1,000} linked to thiomalic acid
3 through an ester bond is tightly cross-linking, as is α,ω -diamino-PEG_{1,000}, through an amide
4 bond.

5
6 The agents may be industrial chemicals or compounds, household chemicals such as cleaners,
7 perfumes or other odorants, deodorants, fertilizers or plant food, foodstuffs such as slow-release
8 food for aquarium fish, therapeutic agents, etc. In a preferred embodiment, the agent is a
9 therapeutically or prophylactically effective agent, generally referred to herein as a therapeutic
10 agent, for controlled release in a bodily compartment of an animal, such as a mammal, preferably
11 a human.

12
13 The therapeutic agents of the invention are not limited to any particular structural type or
14 therapeutic class, and may include small-molecule drugs, peptides and proteins, carbohydrates,
15 and nucleic acids, to name some non-limiting structural compound classes. Small molecule
16 drugs may include, for example, anticancer drugs, cardiovascular drugs, antibiotics, antifungals,
17 antiviral drugs, AIDS drugs such as HIV-1 protease inhibitors and reverse transcriptase
18 inhibitors, antinociceptive (pain) drugs, hormones, vitamins, anti-inflammatory drugs,
19 angiogenesis drugs, and anti-angiogenesis drugs. Among the examples of suitable therapeutic
20 agents are proteins. This includes proteins, peptides, modified proteins and peptides, and
21 conjugates between proteins or peptides and other macromolecules. The protein may be a
22 recombinant protein. For example, candidate agents include erythropoietin, α -interferon, growth
23 hormone and antibodies. Erythropoietin is administered over long periods to promote the

1 formation of red blood cells, such as in conditions including renal failure or cancer
2 therapy-induced anemia. α -Interferon is used to treat certain viral diseases (e.g. hepatitis) and
3 cancers(e.g. hairy cell leukemia). Growth hormone is used for pituitary dwarfism. These
4 compounds are therapeutically effective for certain indications when administered at low doses
5 over an extended period of time, making them good candidates for controlled delivery from a
6 depot administration as described herein, as they otherwise are administered by injection.

7
8 Another group of suitable protein agents are antibodies and antibody fragments, such as those
9 directed against tumor-specific antigens and against inflammatory response proteins such as
10 tumor necrosis factor and interleukin 1, are additional examples of proteins that may be used in
11 the practice of the present invention. As such products usually require frequent parenteral
12 administration, such as by injection, a matrix with an antibody delivered by controlled release
13 provides convenience. The antibody is protected from biodegradative machinery while in the
14 matrix.

15 Another example of a class of therapeutic agents are polysaccharides. Examples include sulfated
16 polysaccharides, such as heparin or calcium spirulan. Heparin is an anticoagulant for which
17 long-term therapy is indicated in various hypercoagulation disorders and for prophylactic use.
18 Chronic anticoagulation therapy is indicated, for example, postoperatively to prevent stroke and
19 pulmonary embolism, and in deep vein thrombosis. Calcium spirulan is a potent antiviral
20 agent against both HIV-1 and HSV-1 (herpes simplex virus) (Hayashi et al., 1996, AIDS
21 Research & Human Retroviruses. 12(15):1463-71).
22
23

1 A further example of suitable therapeutic agents is polynucleotides, such as antisense
2 oligonucleotides. These may be delivered to a particular site within the body using the methods
3 described herein, for sustained delivery to target cells or tissues. Such polynucleotides may be in
4 the form of vectors, gene therapy agents or antisense oligonucleotides. These may be delivered
5 to a particular site within the body using the methods described herein, for sustained delivery to
6 target cells or tissues. Such gene therapy agents include but are not limited to a gene encoding a
7 particular protein or polypeptide domain fragment either as a naked plasmid or introduced in a
8 viral vector. Such vectors include, for example, an attenuated or defective DNA virus, such as
9 but not limited to herpes simplex virus (HSV), papillomavirus, Epstein Barr virus (EBV),
10 adenovirus, adeno-associated virus (AAV), and the like, including retroviral vectors. Defective
11 viruses, which entirely or almost entirely lack viral genes, are preferred. Defective virus is not
12 infective after introduction into a cell. Use of defective viral vectors allows for administration to
13 cells in a specific, localized area, without concern that the vector can infect other cells. Thus, a
14 particular tissue can be specifically targeted.

15 Another example of a therapeutic agent embraced by the invention herein is a vaccine.

16 Administration to an animal of an immunogen in the matrix of the present invention with the
17 proper controlled release kinetics provides the immune system with an antigen for the
18 development of a humoral and/or cellular response. Indeed, fluid flow carrying the released
19 antigen from a subcutaneous depot of the present invention is through lymphatic tissue where
20 the immune response to that antigen may occur.
21

22
23 The foregoing lists and descriptions of therapeutic agents are merely illustrative of examples of

1 various therapeutic agents which may be present singly or in combination in the pharmaceutical
2 compositions of the invention.

3
4 It will be noted that the judicious placement of the matrix of the present invention will permit
5 targeted delivery to a particular site within the body, and furthermore, allow a higher
6 concentration of the therapeutic agent to contact a particular site than achievable if the same
7 therapeutic agent is administered systemically. In particular, administration of an agent which
8 induces apoptosis in dysproliferative conditions, such as a tumor, may be performed by the
9 placement (herein termed administration) of the matrix in the proximity of the tumor, thus
10 delivering the therapeutic agent proximal to the tumor. The same strategy is used for proximal
11 delivery of therapeutic agents to other particular body sites or compartments, such as through the
12 skull into the brain.

13
14 In another embodiment of the present invention, the therapeutic agent may be derivatized to
15 increase its molecular weight, such that it may be better entrapped by and released from the
16 matrix. The derivatization may be, by way of non-limiting example, polymerization or
17 conjugation to poly(ethylene glycol). Such methods of conjugation or polymerization are known
18 to the skilled artisan.

19
20 Alternatively, as described above, the therapeutic agent may be prepared as a suspension of a
21 solid in an aqueous solution of the matrix-forming polymer, thereby becoming entrapped during
22 cross-linking within the matrix in the form of solid particles. Being considerably larger than
23 individual molecules, these solid particles of therapeutic agent will be securely entrapped due to

1 the relatively small pore size of the gels. A given size distribution of the solid particles may be
2 attained by methods known to those skilled in the art. In another embodiment, a carrier
3 molecule, such as human serum albumin (HSA), may be admixed with the therapeutic agent,
4 such as by lyophilizing a solution containing HSA and said therapeutic agent in a preferred ratio
5 of the two components. Thus, the mixture of HSA and therapeutic agent, added in the form of a
6 solid, remains entrapped as a solid during the cross-linking reaction. Dextran, a polysaccharide,
7 may be preferred over HSA as the carrier, since clinical grade dextran of about 70 kDa has a
8 water solubility of about 30 mg/mL, which is >10-fold lower than HSA. Thus, the saturated
9 dextran solution would be less viscous than the HSA solution.

10
11 Since the amount of solid therapeutic agent entrapped is above the solubility limit, then (under
12 ideal conditions) as a given amount of the soluble agent is released from the depot, it is
13 replenished by dissolution of solid therapeutic agent entrapped in the depot. As a result, the
14 concentration of (soluble) therapeutic agent will remain constant, and hence the release rate will
15 remain constant. In the example of quinine sulfate as a therapeutic agent, the water solubility is
16 about 1 mg/mL and about 100 mg of solid quinine sulfate can be used to saturate the polymer
17 solution and then be entrapped within 1 mL of gel. Thus, the highly desired zero order release
18 kinetics should ensue as long as the solution remains saturated, as occurs while the initial 99 mg
19 is being released. Then, only the final 1 mg (1%) of the agent will get released according to first
20 order kinetics, since there is no more solid quinine to replenish the solution. During this tailing
21 off period, the next sustained release dosage of therapeutic agent can be administered to the
22 patient.

1 The possibility of administering the drug as a suspension of solid particles within a subcutaneous
2 gel has an additional advantage with regard to drug loading. For many drugs, the combination of
3 the water solubility of that drug and the amount needed for the duration of the sustained release
4 period would require an unusually large volume of gel. For example, loading 50 mg of quinine
5 sulfate at its solubility limit of 1 mg/mL would require a 50 mL depot. Besides the depot being
6 unsightly, this could make the technology too expensive with regard to cost of polymer and
7 cross-linker. Conversely, the duration of sustained release would have to be kept short to
8 compensate for the limited loading capacity of a poorly water-soluble drug. Yet another
9 consideration is the long-term chemical stability (weeks or months at body temperature) of the
10 therapeutic agent; clearly, said therapeutic agent in most cases would be more stable as a solid
11 rather than as an aqueous solution. Even though some or all of the therapeutic agent is
12 administered as a solid, the present invention also comprises materials and methods for in situ
13 formation of a gel matrix from a mixture containing said polymer(s) and said cross-linking
14 reagent(s). Thus, the invention comprises both the gel and the solid particles of therapeutic agent.
15 Depending on the desired repository site of the matrix of the invention, simply administering a
16 therapeutic agent in the form of solid particles (microparticles, nanoparticles, etc.) could have
17 undesirable attributes. These particles may migrate from the injection site or may be subject to
18 attack by macrophages or soluble degradative enzymes or antibodies, in contrast with the
19 protective environment afforded by a gel. Particles not contained within a gel would not be
20 easily retrievable in case of an adverse side reaction, in contrast with the instant matrix.
21 Furthermore in the present invention, the controlled release kinetics is supported by a small,
22 well-defined gel compartment that can maintain the therapeutic agent as a saturated or
23 near-saturated solution. Moreover, advantageous use of various excipients to maintain the

1 stability of the active agents during residence in the instant compositions will permit the long-
2 term use, and infrequent need to replace, the instant compositions.

3
4 The cross-linked matrix composition of the present invention may be provided in a form such as,
5 but not limited to, a gel, microparticles, and nanoparticles. The composition may be processed
6 for loading into capsules, for example, or for incorporation into another matrix or drug delivery
7 system.

8
9 As mentioned above, release of an entrapped agent may be provided over the course of hours,
10 days, or up to several months. In a further embodiment of the invention in which no release is
11 desired, the compositions of the invention, in particular the cross-linked, emulsion-containing
12 compositions, have their applicability in cosmetic surgery as long-lived, implantable materials to
13 fill in or fill out particular sites in or on the body. In-situ formation of the implant by injection of
14 the components before or during polymerization provides a non-surgical means for placing an
15 inert, shapable mass at any site in the body. In a further embodiment, with the use of reducible
16 cross-links as described above, such a cosmetic implant may be readily removed without surgery
17 after it has achieved its desired purpose. One non-limiting example of such an application is in
18 the theatre, where an actor may desire a temporary altered appearance, such as altered facial
19 features, during the filming or a live production. Post-production, the implant can be
20 depolymerized and flushed or allowed to be absorbed non-surgically.

21
22 In another aspect of the present invention, a method is provided for the controlled release of a
23 therapeutic agent in an animal comprising administration to the animal a therapeutically effective

amount of the therapeutic agent in one of the matrices described above. The matrix may contain more than one therapeutic agent, and an animal may be administered a single therapeutic agent in the form of more than one matrix, each with a particular controlled release kinetic profile.

Administration of the matrix of the present invention is performed to locate the matrix at a desired site for controlled delivery of the therapeutic agent. This may be to a particular body compartment to which the therapeutic agent has a desired targeted effect, or the matrix may be administered to a particular location wherein controlled release may provide the therapeutic agent for distribution throughout the body or to another site from which the administered site drains. Where a number of appropriate sites are possible, one may be selected from which the matrix may be easily removed. The particular site will be determined by the desired effect of the therapeutic agent.

Non-limiting examples of possible sites for administration of the matrix includes subcutaneous, oral, intravenous, intraperitoneal, intradermal, subdermal, intratumor, intraocular, intravisceral, intraglandular, intravaginal, intrasinus, intraventricular, intrathecal, intramuscular, and intrarectal. It will be seen that certain of these sites provides a site from which systemic distribution of the therapeutic agent may occur, for example, intraperitoneal, subcutaneous, and oral. Certain sites may be selected to provide a target tissue or organ to which the therapeutic agent's efficacy is desired, such as intratumor, intravaginal, intraglandular, intrathecal, intraventricular, and intraocular. For example, an antitumor agent may be entrapped in the matrix of the present invention and implanted in or near a tumor, for targeted delivery to the tumor.

1 A subject in whom administration of the pharmaceutical composition of the present invention is
2 preferably a human, but can be any animal. Thus, as can be readily appreciated by one of
3 ordinary skill in the art, the methods and pharmaceutical compositions of the present invention
4 are particularly suited to administration to any animal, particularly a mammal, and including, but
5 by no means limited to, domestic animals, such as feline or canine subjects, farm animals, such
6 as but not limited to bovine, equine, caprine, ovine, and porcine subjects, wild animals (whether
7 in the wild or in a zoological garden), research animals, such as mice, rats, rabbits, goats, sheep,
8 pigs, dogs, cats, etc., avian species, such as chickens, turkeys, songbirds, etc., *i.e.*, for veterinary
9 medical use. In addition, the composition of the present invention may also be used in non-
10 medical situation where controlled release characteristics are desirable, such as, for example,
11 controlled release of fertilizer or anti-parasite agents in the soil near plants; industrial settings,
12 such as purification agents for drinking water tanks, etc. Thus, the term "therapeutic agent" is
13 meant herein to refer to any agent desirous of controlled release.

14 The controlled release of the therapeutic agent from the matrix is believed to occur as a
15 consequence of the diffusion from and/or biodegradation of the matrix by one or more in-vivo
16 degradation pathways. While not wishing to be bound by theory, and by which the inventors
17 herein have no duty to disclose or be bound, it is believed that degradation of the matrix is
18 achieved by local factors at the site of administration such as reducing agents, for example,
19 glutathione; reductases, S-transferases, peptidases, proteases, non-enzymatic hydrolysis,
20 esterases and thioesterases. The varied presence of these various degradation agents in particular
21 compartments in the body provides further guidance on selecting the appropriate site for
22 administration, and also in the preparation of a matrix to provide the desired release kinetics in
23

1 the presence of the particular degradative machinery at the site. Moreover, in compositions of
2 the invention comprising a plurality of phases, the controlled release is further regulated by the
3 presence of, and/or passage through, one or more phases of the composition. For example, as
4 noted above, an insoluble agent in a solid phase may slowly dissolve in the aqueous or oil phase,
5 and this soluble agent then passes out of the composition. A solid phase in the oil phase of an
6 emulsion passes from the solid phase to the oil phase to the aqueous phase. By regulating the
7 properties, relative amounts, presence of excipients, and other parameters of each of the phases,
8 the release characteristics may be adjusted to provide the desired properties for the agent
9 entrapped within the matrix of the compositions of the invention.

10
11 The controlled release characteristics of the pharmaceutical compositions of the invention may
12 be selected for that suited to the particular use. In a preferred embodiment, zero order or pseudo
13 zero order kinetics, i.e., constant release, is desired, where, for example, less than 3% of the
14 therapeutic agent is released from the matrix during the first few hours, and then zero order
15 release continues until at least about 80% of the therapeutic agent is released. First order release
16 kinetics may also be provided.

17
18 In another embodiment of the present invention, the therapeutic agent in the above-described
19 matrix is prepared immediately prior to or during administration to the animal. For example, just
20 prior to administration, a solution, suspension or emulsion containing the therapeutic agent and
21 the polymer can be mixed with a solution containing the cross-linking agent. Upon mixture, the
22 cross-linking of the polymer begins to occur, entrapping the therapeutic agent. As cross-linking
23 proceeds, the mixture changes from a liquid suspension to a gel. The immediately-mixed

1 solutions can be administered as a liquid, for example, by subcutaneous injection, wherein the
2 injected liquid continues to cross-link and change into a matrix at the site of administration. This
3 simplifies the administration of a solid or semi-solid matrix. As the cross-linking traps the
4 therapeutic agent, little is released in a burst during the process.

5
6 In yet another aspect, the present invention is directed to a pharmaceutical composition
7 consisting of a matrix comprising a therapeutic agent exhibiting at least one first controlled
8 release in-vivo kinetic profile, the matrix comprising at least one cross-linked polymer on which
9 at least two thiol groups are present entrapping at least one therapeutic agent. In another
10 embodiment, the therapeutic agent in the aforementioned matrix has at least one second
11 controlled release in-vivo kinetic profile. Controlled release in vivo kinetic profiles refer to the
12 particular release characteristics of the therapeutic agent from the matrix to provide
13 therapeutically effective delivery of the therapeutic agent to the body.

14
15 Another aspect of the invention is the process by which the pharmaceutical compositions of the
16 invention are prepared. The pharmaceutical composition is prepared by cross-linking a polymer
17 on which functional groups are present and are capable of being cross-linked, such as having at
18 least two thiol groups, by any one of various means, in the presence of the therapeutic agent to be
19 entrapped. In a preferred embodiment, at least two thiol groups are present on the polymer.

20 While the following discussion pertains in some instances to the use of thiol-containing
21 polymers, it is understood that in accordance with the general discussions above, that other
22 functional groups on the polymers may be used to achieve similar effects, and the discussions if
23 not dependent on the particular properties of thiol-containing polymers are applicable generally

1 to any and all compositions of the invention.

2
3 In yet a further aspect of the methods and pharmaceutical compositions of the present invention,
4 the polymer or cross-linking agent may additionally comprise a functional group, such as an
5 amino or carboxyl group. The functional group may be derivatized to provide on the polymer or
6 cross-linking agent a moiety such as a label, for example, a contrast/imaging agent, a
7 radionuclide, a chromophore, a fluorophore, a red or near-infrared fluorophore, or a
8 nonradioactive isotope, such that the matrix may be readily located within the body, or the label
9 may be used to monitor degradation of the matrix by detecting a metabolically stable moiety in
10 the urine. The label may be chemically attached to the functional group by, for example,
11 carbodiimide activation or use of a homobifunctional or heterobifunctional cross-linking agent.
12 Examples of contrast/imaging agents include F-19 for MRI, I-126 for X-ray and Tc-99m for
13 radioscintigraphy.

14
15 In a typical example of the preparation of a matrix of the invention, the first step is the synthesis
16 of a polymer on which at least two thiol groups are present. In the case of the amide-linked
17 polymer of α,ω -diamino-PEG with thiomalic acid, the thiomalic acid is first protected as
18 (S-trityl)-thiomalic acid, as follows. Equimolar quantities of α,ω -diamino-PEG (MW 3,400,
19 Shearwater Polymers) and (S-trityl)-thiomalic acid were dissolved in methylene chloride, and 3.5
20 equivalents of 1,3-diisopropylcarbodiimide (DIPC, Aldrich) was added to carry out a direct
21 polycondensation at room temperature with 0.5 equivalent of 4-(dimethylamino)-pyridine
22 (DMAP, Aldrich) and p-toluenesulfonic acid monohydrate (PTSA, Aldrich) as catalysts. The
23 reaction mixture was precipitated with cold ethyl ether to obtain a white polymer product, which

1 was treated with 100% trifluoroacetic acid for 2 hours to remove the protecting trityl groups from
2 the polymer. The deprotected polymer was precipitated in cold ethyl ether, washed 5 times with
3 ether and dried under vacuum. The molecular weight of the resulting PEG-thiomalic acid
4 polymer was measured by size exclusion chromatography, using PEGs of defined molecular
5 weights (Shearwater Polymers) for calibration of the column.

6
7 These polymers are then to be used for the preparation of the matrix entrapping the therapeutic
8 agent in the presence of two or more phases. As mentioned above, use of polymers made from
9 the smaller PEG subunits would result in a matrix having more closely spaced cross-links,
10 resulting in a slower rate of diffusion of entrapped therapeutic agents, especially higher
11 molecular weights, out of the matrix. Amide bonds, resulting from use of the diamino-PEGs, are
12 expected to be considerably more stable *in vivo* than are ester bonds, which corresponds to a
13 lower rate of degradation of the matrix *in vivo*.

14
15 For matrix formation, a preferred cross-linking reagent is α,ω -divinylsulfone-PEG (Shearwater
16 Polymers). The vinylsulfone functional group reacts readily and specifically with thiol groups on
17 the matrix-forming polymer, but will not react with disulfide bonds, such as present in a protein
18 with disulfide bonds. As mentioned above, the possibility of cleavage of any disulfide bond in
19 the therapeutic agent can be minimized or essentially prevented by providing steric hindrance to
20 the thiol groups in the thiol-containing polymer.

21
22 Another factor influencing the release rate of the therapeutic agent is the size and the shape of the
23 matrix depot. The greater the ratio of surface area to volume, the shorter the duration of release.

1 For example, a sheet-like depot would be expected to release the encapsulated agent much faster
2 than would a spherical depot of the same mass. One large sphere would release the agent more
3 slowly than would many small spheres of the same total mass. The selection of the size and
4 shape of the matrix will be readily determinable by a skilled artisan based on desired
5 characteristics of release of the particular therapeutic agent. Other factors include the relative
6 amount of each of the multiple phases present in the composition, the phase(s) in which the
7 active agent or agents is or are present, the solubility of the agent(s) and partitioning between the
8 phases, etc. By using the teaching herein, the skilled artisan can readily determine for a
9 particular use and agent(s) the proper features of the desired composition and the means to
10 prepare it.

11
12
13
14
15
16 As mentioned above, the matrix may be administered just after mixing the polymer on which at
17 least two thiol groups are present with the cross-linking agent, in the presence of the therapeutic
18 agent, in the one or more phases, such that the mixture may be injected in liquid form but the
19 matrix solidifies into the cross-linked form soon thereafter. For example, to practice this aspect
20 of the invention, a dual-syringe pump may be used for making and administering the mixture.
21 For example, one syringe will be filled with 0.5 mL of matrix-forming polymer and the
22 therapeutic agent in a plurality of phases, while the other syringe will be filled with 0.5 mL of the
23 cross-linker solution (or the therapeutic agent may be mixed in this syringe), both at the optimal
concentrations for the cross-linking reaction. The concentrations selected for these two solutions
will be that appropriate to create the matrix with the appropriate controlled release kinetic profile.
The pump will be set at a constant flow rate (e.g. 0.1 mL/min). The two solutions will be mixed
in a tee-fitting and the mixture will be injected. The mixture becomes viscous as it flows through

1 teflon tubing for a specified time. The mixed solution may be injected to the site of
2 administration, whereupon the solution polymerizes into a multiphase hydrogel matrix. More
3 simply, all components may be mixed just prior to administration.

4
5 More simply, all components can be mixed in one syringe just prior to administration. The rate
6 at which the gel forms by the cross-linking reaction is preferably in a time frame of a few
7 minutes. This rate may be controlled by the type of functional group on the cross-linking reagent
8 and by the pH of the reaction, being slower at pH 6 compared with pH 7.

9
10
11
12
13
14
15
16 With regard to the administration of the matrix as described above, in one embodiment, may
17 comprise, for example, several injections of 1 microliter each, perhaps repeated at multiple sites
18 around the body, whereby the number and volume of the injections corresponds to a particular
19 pharmacokinetic profile. As noted above, the fluid would be a partially cross-linked viscous
20 matrix as it enters the skin, thereby already entrapping the drug. Microparticles, perhaps
21 uniformly sized at 1 cubic millimeter (about 1 microliter), would harden within minutes as the
22 cross-linking reaction goes to completion. Alternatively, a single needle injection may be used to
23 produce a subcutaneous depot that may be easier to remove surgically in case of an adverse
24 reaction to the depot or the drug.

25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
217

1 therapeutic agent, the longevity of the matrix, among other factors. Each therapeutic agent will
2 require a particular set of factors to provide the matrix with the correct profile for therapeutic use.
3 In particular, the molecular weight and physical interaction between the agent and the polymers
4 comprising the matrix will participate in the profile. For the practice of the invention, a
5 particular set of preparation and operating conditions will be established for each therapeutic
6 agent and, in more particular, the desired controlled release profile for that agent. It is well
7 within the realm of the skilled artisan, based on the teaching herein, to determine the matrix
8 components and other factors in the preparation of a suitable range of conditions for preparing a
9 matrix for a particular therapeutic agent which exhibits the desired profile.

10
11 Further to the typical procedure described above for the preparation of the matrix of the present
12 invention, variables for the protein solution include but are not limited to protein concentration,
13 pH, salt content and presence of other excipients and stabilizers. The protein may be modified,
14 such as by pegylation, to increase its size and, thereby, decrease its release rate.

15
16 In a further aspect of the present invention, a particular release rate may be achieved using a
17 mixture of two or more starting polymer subunits to prepare the thiol-containing polymer or
18 using a mixture of two or more polymers during the cross-linking/entrapment process. A delayed
19 release product may be prepared by first entrapping the protein using an ester-type
20 polymer, followed by coating or encapsulating these resulting particles using an amide-type
21 polymer. The desired release kinetics for the final product may be achieved by administering to
22 the patient a blend of two or more differently and separately cross-linked, entrapped protein
23 preparations. Other means for making a product with a desired release profile will be apparent to

1 the skilled artisan based on the teachings herein and should be considered to be within the scope
2 and spirit of the present invention. As mentioned above, for any particular matrix, the release rate
3 must be determined empirically in vivo, since it is dependent on many factors, including the size
4 of the protein, diffusion from the matrix and the rate of degradation of the cross-linked polymer
5 matrix due to the action of esterases, peptidases and reducing agents at the site of the depot.

6
7 The present invention may be better understood by reference to the following non-limiting
8 Examples, which are provided as exemplary of the invention. The following examples are
9 presented in order to more fully illustrate the preferred embodiments of the invention. They
10 should in no way be construed, however, as limiting the broad scope of the invention.

11 12 13 14 15 16 17 18 19 20 21 22 23 **Example 1**

Entrapment of quinine sulfate monohydrate in a thiol containing polymer hydrogel through a suspension system.

17 A thiol-containing polymer was prepared from thiomalic acid and α,ω -diamino-PEG as follows.
18 One equivalent of thiomalic acid and 3 equivalents of trityl chloride were dissolved in
19 dimethylformamide (DMF). The reaction was carried out at room temperature with stirring
20 overnight. The reaction mixture was loaded onto a silica gel column and the eluted fractions
21 containing trityl-thiomalic acid were collected and evaporated to dryness. Equimolar quantities
22 of α,ω -diamino-PEG (MW 3,400; Shearwater Polymers, Inc. Huntsville, AL) and thiol
23 group-protected thiomalic acid as prepared above were dissolved in methylene chloride, and 3.5

equivalents of 1,3-diisopropylcarbodiimide (DIPC, Aldrich, Milwaukee, WI) were added to carry out a direct polycondensation at room temperature with 0.5 equivalent of 4-(dimethylamino)-pyridine (DMAP, Aldrich, Milwaukee, WI) and p-toluenesulfonic acid monohydrate (PTSA, Aldrich, Milwaukee, WI) as catalyst. The reaction mixture was precipitated with cold ethyl ether to obtain a white polymer product. The polymer was treated with 100% trifluoroacetic acid (TFA) for 2 hours to remove the protecting trityl groups from the polymer pendant chain. The deprotected polymer was precipitated in cold ethyl ether, washed 5 times with ether and dried under vacuum.

Sixteen mg of the foregoing polymer was dissolved in 300 microliters of PBS, pH 7.4. Fifty mg of quinine sulfate monohydrate (Aldrich Chemical Co., Milwaukee, WI) was added into the polymer solution to form a suspension. Then, 4.7 mg PEG-(VS)₂ (MW 2000 Da, Shearwater Polymers, Inc., Huntsville, AL) was dissolved in 100 microliters of PBS, pH 7.4. The two solutions were mixed thoroughly in a 1.5 mL Eppendorf tube. The mixture was allowed to stand at room temperature (25 degree C) until the hydrogel formed ("DepoGel formulation I").

Example 2

Entrapment of quinine sulfate monohydrate in a thiol containing polymer hydrogel through an emulsion system.

A thiol-containing polymer prepared from α,ω -dihydroxy-PEG and thiomalic acid was prepared as follows. Equimolar quantities of α,ω -dihydroxy-PEG and thiomalic acid were dissolved in methylene chloride, and 3.5 equivalent of 1,3-diisopropylcarbodiimide (DIPC, Aldrich,

1 Milwaukee, WI) were added to carry out a direct polycondensation at room temperature with 0.5
2 equivalent of 4-(dimethylamino)-pyridine (DMAP, Aldrich, Milwaukee, WI) and
3 p-toluenesulfonic acid monohydrate (PTSA, Aldrich, Milwaukee, WI) as catalyst. The reaction
4 mixture was precipitated with cold ethyl ether to obtain a thiol-containing polymer.

5
6 Sixteen mg of the foregoing thiol-containing polymer was dissolved in 200 microliters of PBS,
7 pH 7.4. To this, 200 microliters of ethyl myristate (Aldrich Chemical Co., Milwaukee, WI) was
8 added as the oil phase and 24 mg of sodium dodecylsulfate (Bio-Rad, Hercules, CA) as the
emulsifier. The mixture was mixed thoroughly to form an emulsion system. Fifty mg of quinine
sulfate monohydrate (Aldrich Chemical Co., Milwaukee, WI) was added into the above emulsion
system. Then, 4.7 mg PEG-(VS)₂ (MW 2000 Da, Shearwater Polymers, Inc., Huntsville, AL)
was dissolved in 100 mL of PBS, pH 7.4. After thorough mixing in a 1.5 mL Eppendorf tube,
the mixture was allowed to stand at room temperature (25 degree C) until the hydrogel formed
("DepoGel formulation II").

16 Example 3

17 Release of quinine sulfate monohydrate from thiol containing polymer hydrogels.

18
19 To conduct a release study, 2 mL of PBS, pH 7.4 was added to the polymer hydrogel in a 5 mL
20 test tube and allowed to incubate the hydrogel at room temperature for pre-selected time periods
21 with rotation (about 100 rpm). The supernatant from the hydrogel was removed for fluorescence
22 measurement and to add fresh PBS for the next incubation.

1 Fluorescence measurements of released quinine sulfate monohydrate were performed using
2 FALCON microtiter plates from Becton Dickinson (Lincoln Park, NJ) on a CytoFluorä II
3 fluorescence multi-well plate reader (PerSeptive Biosystems, Framingham, MA). For each
4 measurement, 100 microliters of release sample is mixed with 100 microliters of 1 M sulfuric
5 acid in a well of the microtiter plate. An excitation wavelength of 360 nm and an emission
6 wavelength of 460 nm are used for fluorescence measurements. Based on fluorescence
7 measurement of each collected release sample, the release profiles of quinine sulfate
8 monohydrate from thiol containing polymer hydrogels in PBS, pH 7.4 at 25 degree C are shown
in Figure 1. DepoGel formulation I shows the quinine release from a thiol containing polymer
hydrogel through a suspension system. DepoGel formulation II shows the quinine release from a
thiol containing polymer hydrogel through a emulsion system.

Example 4

Further example of the in-vitro release of quinine sulfate

16
17 Single-phase system (R-gel): In a test tube, 16 mg of Thiol-PEG polymer were dissolved in 400
18 µl of PBS (pH=7.4), 50 mg of quinine sulfate was added to the polymer solution to form a
19 suspension. In another test tube, 4.7 mg of PEG-divinylsulfone (PEGDVS) were dissolved in 100
20 µl of PBS (pH 7.4) as the cross-linker solution. The cross-linker solution is added into the above
21 prepared suspension and it is mixed thoroughly. A polymer hydrogel is formed in about 3
22 minutes.
23

Two-phase (emulsion) system (E-gel): In a test tube, 16 mg of Thiol-PEG polymer and 24 mg SDS are dissolved in 200 μ l of PBS (pH=7.4). 200 μ l of ethyl myristate ("Oil") are added and mixed thoroughly to form an emulsion. 50 mg of quinine sulfate is added to above emulsion. In another test tube, 4.7 mg of PEG-divinylsulfone (PEGDVS) are dissolved in 100 μ l of PBS (pH 7.4) as the cross-linker solution. The cross-linker solution is added into the above prepared emulsion and mixed thoroughly. A polymer hydrogel is formed in about 3 minutes.

Release conditions and Sample collection: To each test tube containing R-gel and E-gel, 2 mL of PBS is added. The test tubes are set on a rotational shaker (250 rpm) at room temperature (25 °C). At pre-selected time points, all solution is removed for sample analysis and 2 mL of fresh PBS is added into each tube.

Sample analysis: For each collected sample, 100 μ l of sample is mixed with 100 μ l of 1 M H_2SO_4 solution in a microplate. Fluorescence measurements were performed on a CytoFluor(tm) II fluorescence multi-well plate reader (PerSeptive Biosystems, Framingham, MA).

Results: Fig. 2 shows that the in-vitro release rate of a small molecule drug, quinine sulfate from E-gel (with a lipid excipient) which displays an apparent zero-order release profile and is much slower than release without excipient (R-gel). Only 40% of the quinine is released from the E-gel with excipient in 4 months of study, whereas essentially 100% is released without excipient in 2 months from the R-gel. There is no substantial initial burst effect.

Example 5

In-vivo release study of quinine sulfate

Animal model: New Zealand great white rabbits were used for an in-vivo release study. The average weight of the rabbits was 3.0 kg. Three groups of rabbits were used for the study and each group contains 3 rabbits. Group A was used for subcutaneous injection of quinine sulfate solution (not in a polymer system). Group B was used for subcutaneous injection of quinine sulfate in Rgel, as described above. Group C was used for subcutaneous injection of quinine sulfate in Egel, the lipid emulsion described above.

Preparation of plain injection of quinine sulfate: For plain injection, 1 mL of 1 mg/mL quinine sulfate solution was injected subcutaneously into the upper back area of each rabbit in Group A.

Preparation of Rgel: For Rgel preparation, 16 mg of Thiol-PEG polymer is dissolved in 400 μ l of PBS (pH=7.4), and 50 mg of quinine sulfate is added to the polymer solution to form a suspension. 4.7 mg of cross-linker, PEGDVS is dissolved in 100 μ l of PBS (pH 7.4). The cross-linker solution is drawn into a 1 mL syringe first, then draw the thiol-PEG polymer solution into the same syringe. It was mixed thoroughly by drawing up and push down the syringe plunger several times. The solution gradually became viscous in 2 minutes; then this viscous solution was administered subcutaneously into the upper back area of each rabbit in Group B.

Preparation of Egel: For Egel preparation, 16 mg of Thiol-PEG polymer and 24 mg SDS were dissolved in 200 μ l of PBS (pH=7.4). 200 μ l of ethyl myristate ("Oil") was added and mixed thoroughly to form an emulsion. 50 mg of quinine sulfate was added to above emulsion. In another test tube, 4.7 mg of cross-linker, PEGDVS was dissolved in 100 μ l of PBS (pH 7.4). The cross-linker solution is drawn into a 1 mL syringe, followed by the thiol-PEG polymer solution. The syringe contents were mixed thoroughly by drawing the syringe plunger up and down several times. The solution gradually became viscous in 2 minutes; the viscous solution was administered subcutaneously into the upper back area of each rabbit in Group C.

Sample collection: At pre-selected time points, 2 mL of blood was collected from vein of the rabbit ear into an EDTA-treated test tube. The blood was centrifuged at 3000x g at 4 °C to obtain about 1 mL of plasma. All plasma samples were kept at -70 °C until analysis.

Sample analysis: Reverse-phase HPLC method was used for plasma sample analysis under following conditions: HPLC column: Princeton SPHER ULTIMA C18 100 Å 5 μ 150x4.6 nm Mobile Phase: 95/5 25 mM KH₂PO₄, pH3/Methanol. Flow rate: 1 mL/min

Sample treatment: Plasma was precipitated with 2 volumes of cold methanol, vortexed and centrifuges at 1500 x g for 10 min. The supernatant (10 μ l) was injected into the HPLC column.

Results: Fig. 3 shows the in-vivo release of a small molecule drug, quinine sulfate from polymer systems of the present invention in rabbits. Egel containing a lipid excipient displays a slower

1 drug release than that of Rgel, the polymer system without excipient. There are no substantial
2 initial burst effects in either case.

4 **Example 6**

5 **In-vitro release study of salmon calcitonin**

6
7 Preparation of polymer system without lipid excipient (Rgel): In a series of test tubes, 20 mg of
8 soluble polymer was dissolved in PBS (pH 5.5) to yield 400 μ l of solution in each tube. In
9 another series of test tubes, 1 mg of cross-linker and 1 mg of salmon calcitonin were dissolved in
10 100 μ l of PBS (pH 5.5). The two solutions were mixed thoroughly at room temperature (25 $^{\circ}$ C),
11 and a series of polymer hydrogels was formed in about 1 min.

12 Preparation of polymer system containing lipid excipient (Egel): In a series of test tubes, 20 mg
13 of soluble polymer and varying amounts of lipid excipient were dissolved in PBS (pH 5.5) to
14 yield 400 μ l of solution in each tube. In another series of test tubes, 1 mg of cross-linker and 1
15 mg of salmon calcitonin were dissolved in 100 μ l of PBS (pH 5.5). The two solutions were
16 mixed thoroughly at room temperature (25 $^{\circ}$ C), and a series of polymer hydrogels was formed in
17 about 1 min.
18

19
20 Release Conditions and Sample Collection: To each test tube containing a polymer system, 1
21 mL of PBS (pH 5.5) was added. The test tubes were set on a rotational shaker (300 rpm) at room
22 temperature (25 $^{\circ}$ C). At pre-selected time points, all solution was removed from each tube for
23 sample analysis and 1 mL of fresh PBS (pH 5.5) was added to each tube.

1 Sample analysis: All collected samples are analyzed by HPLC as described above.

2
3 Results: Fig. 4 shows that the use of the lipid excipient-containing cross-linked polymer
4 significantly slowed the in-vitro release rate of a peptide drug, salmon calcitonin (sCT) from the
5 polymer.

7 **Example 7**

8 **In-vivo release study of salmon calcitonin**

9
10 Animal model: New Zealand great white rabbits were used for an in-vivo release study. The
11 average weight of the rabbits was 3.0 kg. Two groups of rabbits were used for the study and each
12 group contained 3 rabbits. Group A was used for subcutaneous injection of salmon calcitonin in
13 Rgel. Group B was used for subcutaneous injection of salmon calcitonin in Egel.

14
15 Preparation of Rgel (also referred to herein as Depogel Formulation I): For Rgel preparation, 40
16 mg of Thiol-PEG polymer is dissolved in 800 μ l of PBS (pH=7.4). 10 mg of salmon calcitonin
17 and 2 mg of the cross-linker 1,11-bis-maleimidotetraethylene glycol [BM(EG)₄] were dissolved
18 in 200 μ l of PBS (pH 7.4). The cross-linker solution was drawn into a 3 mL syringe first, then
19 the thiol-PEG polymer solution was drawn into the same syringe. It was mixed thoroughly by
20 drawing up and pushing down the syringe plunger several times. The solution gradually became
21 viscous within 1 minute; then this viscous solution is administered subcutaneously into the upper
22 back area of each rabbit in Group A. A soft, round-shaped depot is formed at the injection site
23 upon injection.

Preparation of Egel (also referred to herein as Depogel Formulation II): For Egel preparation, 40 mg of Thiol-PEG polymer and 5 mg SDS are dissolved in 700 μ l of PBS (pH=7.4). 100 μ l of ethyl myristate ("Oil") was added and mixed thoroughly to form an emulsion. In another test tube, 10 mg of salmon calcitonin and 2 mg of cross-linker, BM(EG)₄ were dissolved in 200 μ l of PBS (pH 7.4). The cross-linker solution was drawn into a 3 mL syringe, followed by the thiol-PEG polymer solution containing oil droplets (i.e. an emulsion system). The syringe contents were mixed thoroughly by drawing the syringe plunger up and down several times. The solution gradually became viscous within 1 minute; the viscous solution was administered subcutaneously into the upper back area of each rabbit in Group B. A soft, round-shaped depot was formed at the injection site upon injection.

Sample collection: At pre-selected time points, 1 mL of blood was collected from vein of the rabbit ear into a heparin-treated test tube. The blood was centrifuged at 3000g at 4 °C to obtain about 0.5 mL of plasma. All plasma samples were kept at -70 °C until analysis.

Sample analysis: A radioimmunoassay (RIA) was used for plasma sample analysis to determine the salmon calcitonin level in rabbit plasma.

Results: Fig. 5 shows the in-vivo release of a peptide drug, salmon calcitonin, from polymer systems of the invention in rabbits. Egel, containing a lipid excipient, displays a slower drug release than that of Rgel, without excipient. There are no substantial initial burst effects in either case.

1
2 The present invention is not to be limited in scope by the specific embodiments described herein.
3 Indeed, various modifications of the invention in addition to those described herein will become
4 apparent to those skilled in the art from the foregoing description and the accompanying figures.
5 Such modifications are intended to fall within the scope of the appended claims.

6
7 Various publications are cited herein, the disclosures of which are incorporated by reference in
8 their entireties.

09/03/2012 10:13:01